



# Medical Assistance BULLETIN

<b>ISSUE DATE</b> November 14, 2025	<b>EFFECTIVE DATE</b> January 5, 2026	<b>NUMBER</b> *See below
<b>SUBJECT</b>  Prior Authorization of Lipotropics, Other – Pharmacy Services		<b>BY</b>  Sally Kozak Deputy Secretary Office of Medical Assistance Programs

**IMPORTANT REMINDER:** All providers must revalidate the Medical Assistance (MA) enrollment of each service location every 5 years. Providers should log into PROMISe to check the revalidation dates of each service location and submit revalidation applications at least 60 days prior to the revalidation dates. Enrollment (revalidation) applications may be found at: <https://www.pa.gov/en/agencies/dhs/resources/for-providers/provider-enrollment-information/provider-enrollment-documents.html>.

**PURPOSE:**

The purpose of this bulletin is to issue updated handbook pages that include the requirements for prior authorization and the type of information needed to evaluate the medical necessity of prescriptions for Lipotropics, Other submitted for prior authorization.

**SCOPE:**

This bulletin applies to all licensed pharmacies and prescribers enrolled in the Medical Assistance (MA) Program. The guidelines to determine the medical necessity of Lipotropics, Other will be utilized in the fee-for-service and managed care delivery systems. Providers rendering services to MA beneficiaries in the managed care delivery system should address any questions related to the prior authorization of Lipotropics, Other to the appropriate managed care organization.

**BACKGROUND:**

The Department of Human Services’ (Department) Pharmacy and Therapeutics (P&T) Committee reviews published peer-reviewed medical literature and recommends the following:

*01-26-25	09-26-25	27-26-25	33-26-25
02-26-25	11-26-25	30-26-25	
03-26-25	14-26-25	31-26-25	
08-26-25	24-26-25	32-26-25	

**COMMENTS AND QUESTIONS REGARDING THIS BULLETIN SHOULD BE DIRECTED TO:**

Fee-for-Service Provider Service Center: 1-800-537-8862

Visit the Office of Medical Assistance Programs website at:  
<https://www.pa.gov/en/agencies/dhs/departments-offices/omap-info.html>

- Preferred or non-preferred status for new drugs and products in therapeutic classes already included on the Statewide Preferred Drug List (PDL).
- Changes to the statuses of drugs and products on the Statewide PDL from preferred to non-preferred and non-preferred to preferred.
- Therapeutic classes of drugs and products to be added to or deleted from the Statewide PDL.
- New quantity limits.
- New guidelines or revisions to existing guidelines to evaluate the medical necessity of prescriptions submitted for prior authorization.

### **DISCUSSION:**

During the September 25, 2025, meeting, the P&T Committee recommended revisions to the medical necessity guidelines for Lipotropics, Other to add a requirement for prior authorization and corresponding medical necessity guidelines for *APOC-III*-directed antisense oligonucleotides and clarify the guidelines related to angiotensin-like 3 inhibitors and microsomal triglyceride transfer protein inhibitors.

The revisions to the guidelines to determine medical necessity of prescriptions for Lipotropics, Other submitted for prior authorization, as recommended by the P&T Committee, were subject to public review and comment and subsequently approved for implementation by the Department.

### **PROCEDURE:**

The procedures for prescribers to request prior authorization of Lipotropics, Other are located in SECTION I of the Prior Authorization of Pharmaceutical Services Handbook. The Department will take into account the elements specified in the clinical review guidelines (which are included in the provider handbook pages in the SECTION II chapter related to Lipotropics, Other) when reviewing the prior authorization request to determine medical necessity.

As set forth in 55 Pa. Code § 1101.67(a), the procedures described in the handbook pages must be followed to ensure appropriate and timely processing of prior authorization requests for drugs and products that require prior authorization.

### **ATTACHMENTS:**

Prior Authorization of Pharmaceutical Services Handbook - Updated pages

### **RESOURCES:**

Prior Authorization of Pharmaceutical Services Handbook – SECTION I  
Pharmacy Prior Authorization General Requirements

<https://www.pa.gov/en/agencies/dhs/resources/pharmacy-services/pharmacy-prior-authorization-general-requirements.html>

Prior Authorization of Pharmaceutical Services Handbook – SECTION II  
Pharmacy Prior Authorization Guidelines

<https://www.pa.gov/en/agencies/dhs/resources/pharmacy-services/clinical-guidelines.html>

MEDICAL ASSISTANCE HANDBOOK  
PRIOR AUTHORIZATION OF PHARMACEUTICAL SERVICES

**I. Requirements for Prior Authorization of Lipotropics, Other**

**A. Prescriptions That Require Prior Authorization**

Prescriptions for Lipotropics, Other that meet any of the following conditions must be prior authorized:

1. A non-preferred Lipotropics, Other. See the Preferred Drug List (PDL) for the list of preferred Lipotropics, Other at: <https://papdl.com/preferred-drug-list>.
2. A Lipotropics, Other with a prescribed quantity that exceeds the quantity limit. The list of drugs that are subject to quantity limits, with accompanying quantity limits, is available at: <https://www.pa.gov/en/agencies/dhs/resources/pharmacy-services/quantity-limits-daily-dose-limits.html>.
3. A proprotein convertase subtilisin/kexin type 9 (PCSK9) inhibitor (e.g., alirocumab, evolocumab, inclisiran).
4. An adenosine triphosphate-citrate lyase (ACL) inhibitor (e.g., bempedoic acid, bempedoic acid/ezetimibe).
5. A microsomal triglyceride transfer protein (MTP) inhibitor (e.g., lomitapide).
6. An angiopoietin-like 3 (ANGPTL3) inhibitor (e.g., evinacumab).
7. An *APOC-III*-directed antisense oligonucleotide (ASO) (e.g., olezarsen).

**B. Review of Documentation for Medical Necessity**

In evaluating a request for prior authorization of a prescription for a Lipotropics, Other, the determination of whether the requested prescription is medically necessary will take into account whether the beneficiary:

1. Is prescribed the requested Lipotropics, Other for the treatment of a diagnosis that is indicated in the U.S. Food and Drug Administration (FDA)-approved package labeling or a medically accepted indication; **AND**
2. Is prescribed a dose that is consistent with FDA-approved package labeling, nationally recognized compendia, or peer-reviewed medical literature; **AND**
3. Is age-appropriate according to FDA-approved package labeling, nationally recognized compendia, or peer-reviewed medical literature; **AND**
4. Does not have a contraindication to the prescribed drug; **AND**

MEDICAL ASSISTANCE HANDBOOK  
PRIOR AUTHORIZATION OF PHARMACEUTICAL SERVICES

5. For treatment of a lipid disorder, has documentation of results of a lipid profile within three months prior to the request for the Lipotropics, Other; **AND**
6. For a PCSK9 inhibitor, **all** of the following:
  - a. Has a history of **one** of the following:
    - i. Failure to achieve goal LDL-C or percentage reduction of LDL-C while adherent to treatment with the maximum tolerated dose of a high-intensity statin for greater than or equal to three months,
    - ii. **Both** of the following:
      - a) A temporally related intolerance<sup>1</sup> to two high-intensity statins that occurred after **both** of the following:
        - (i) Modifiable comorbid conditions that may enhance statin intolerance were ruled out and/or addressed by the prescriber as clinically indicated (e.g., hypothyroidism, vitamin D deficiency)
        - (ii) All possible drug interactions with statins were addressed by **all** of the following (if clinically appropriate):
          - a. Dose decrease of the interacting non-statin drug,
          - b. Discontinuation of the interacting non-statin drug,
          - c. Change to an alternative statin that has a lower incidence of drug interactions
      - b) **One** of the following:
        - (i) Therapeutic failure while adherent to treatment for greater than or equal to three consecutive months with the lowest FDA-approved daily dose or alternate-day dosing of any statin
        - (ii) A temporally related intolerance to the lowest FDA-approved daily dose or alternate-day dosing of any statin,
    - iii. A contraindication to statins,
  - b. Has **one** of the following:

---

<sup>1</sup> Temporally related intolerance of a statin is defined as the occurrence of symptoms and/or lab abnormalities upon initiation of a statin, resolution of symptoms and/or lab abnormalities upon discontinuation of a statin, and recurrence of symptoms and/or lab abnormalities after rechallenge with the same statin at the same dose.

MEDICAL ASSISTANCE HANDBOOK  
PRIOR AUTHORIZATION OF PHARMACEUTICAL SERVICES

- i. A history of therapeutic failure of while adherent to treatment with ezetimibe in combination with the maximum tolerated dose of the highest-tolerated intensity statin (if clinically appropriate) for greater than or equal to three consecutive months,
  - ii. A contraindication or an intolerance to ezetimibe,
  - iii. An LDL-C that is greater than 25% above goal LDL-C while adherent to treatment with the maximum tolerated dose of the highest-tolerated intensity statin (if clinically appropriate) for greater than or equal to three consecutive months,
- c. Is prescribed the requested PCSK9 inhibitor in addition to **one** of the following:
- i. For treatment of homozygous familial hypercholesterolemia (HoFH), standard lipid-lowering treatments as recommended by current consensus guidelines<sup>2</sup>
  - ii. For treatment of all other conditions, the maximum tolerated dose of the highest-tolerated intensity statin (if clinically appropriate),
- d. If currently using a different PCSK9 inhibitor, will discontinue use of that PCSK9 inhibitor prior to starting the requested PCSK9 inhibitor,
- e. For a non-preferred PCSK9 inhibitor, has **one** of the following:
- i. A history of therapeutic failure of at least one preferred PCSK9 inhibitor approved or medically accepted for the beneficiary's diagnosis
  - ii. A contraindication or an intolerance to the preferred PCSK9 inhibitors approved or medically accepted for the beneficiary's diagnosis;

**AND**

7. For an ACL inhibitor, **all** of the following:
- a. Has a history of **one** of the following:
    - i. Failure to achieve goal LDL-C or percentage reduction of LDL-C while adherent to treatment with the maximum tolerated dose of a high-intensity statin for greater than or equal to three months,
    - ii. **Both** of the following:
      - a) A temporally related intolerance to two high-intensity statins that occurred after **both** of the following:
        - (i) Modifiable comorbid conditions that may enhance statin intolerance were ruled out and/or addressed by the prescriber as clinically indicated (e.g.,

---

<sup>2</sup> e.g., American Heart Association/American College of Cardiology, American Association of Clinical Endocrinologists/American College of Endocrinology, American Diabetes Association, National Lipid Association, European Society of Cardiology/European Atherosclerosis Society, International Familial Hypercholesterolaemia Foundation, International Atherosclerosis Society

MEDICAL ASSISTANCE HANDBOOK  
PRIOR AUTHORIZATION OF PHARMACEUTICAL SERVICES

hypothyroidism, vitamin D deficiency)

(ii) All possible drug interactions with statins were addressed by **all** of the following (if clinically appropriate):

- a. Dose decrease of the interacting non-statin drug,
- b. Discontinuation of the interacting non-statin drug,
- c. Change to an alternative statin that has a lower incidence of drug interactions

b) **One** of the following:

- (i) Therapeutic failure while adherent to treatment for greater than or equal to three consecutive months with the lowest FDA-approved daily dose or alternate-day dosing of any statin
- (ii) A temporally related intolerance to the lowest FDA-approved daily dose or alternate-day dosing of any statin,

iii. A contraindication to statins,

b. Has **one** of the following:

- i. A history of therapeutic failure of while adherent to treatment with ezetimibe in combination with the maximum tolerated dose of the highest-tolerated intensity statin (if clinically appropriate) for greater than or equal to three consecutive months
- ii. A contraindication or an intolerance to ezetimibe,

c. Is prescribed the requested ACL inhibitor in addition to the maximum tolerated dose of the highest-tolerated intensity statin (if clinically appropriate),

d. If currently taking simvastatin or pravastatin, will not be using the requested ACL inhibitor concomitantly with simvastatin at a dose of greater than 20 mg daily or pravastatin at a dose of greater than 40 mg daily;

**AND**

8. For an ANGPTL3 inhibitor or MTP inhibitor, **all** of the following:

a. Is prescribed the requested drug by or in consultation with a cardiologist, endocrinologist, or other provider specializing in lipid disorders,

b. For treatment of HoFH, has a diagnosis of HoFH in accordance with current consensus guidelines,

c. **One** of the following:

MEDICAL ASSISTANCE HANDBOOK  
PRIOR AUTHORIZATION OF PHARMACEUTICAL SERVICES

- i. Has a history of therapeutic failure of or a contraindication or an intolerance to PCSK9 inhibitors
  - ii. Has results of genetic testing that are positive for mutations associated with lack of response to PCSK9 inhibitors,
- d. Is prescribed the requested drug in addition to standard lipid-lowering treatments as recommended by current consensus guidelines;

**AND**

9. For icosapent ethyl, **all** of the following:
- a. **One** of the following:
    - i. Has a history of clinical ASCVD,
    - ii. **Both** of the following:
      - a) Has diabetes mellitus
      - b) Has two additional ASCVD risk factors (e.g., 50 years of age or older, cigarette smoking, hypertension, HDL-C less than or equal to 40 mg/dL for males or 50 mg/dL for females, hs-CRP greater than 3.00 mg/L, CrCl less than 60 mL/min, retinopathy, micro- or macroalbuminuria, ABI less than 0.9]),
    - iii. Has a history of therapeutic failure of or a contraindication or an intolerance to the preferred Lipotropics, Other approved or medically accepted for the beneficiary's diagnosis,
  - b. Has fasting triglycerides greater than or equal to 150 mg/dL,
  - c. Has **one** of the following:
    - i. A history of therapeutic failure of while adherent to treatment with maximum tolerated doses of two different statins for greater than or equal to three consecutive months each,
    - ii. A history of statin intolerance after modifiable risk factors have been addressed,
    - iii. A contraindication to statins;

**AND**

10. For an *APOC-III*-directed ASO, **both** of the following:
- a. Is prescribed the requested drug by or in consultation with a cardiologist, endocrinologist, gastroenterologist, or other provider specializing in lipid disorders

MEDICAL ASSISTANCE HANDBOOK  
PRIOR AUTHORIZATION OF PHARMACEUTICAL SERVICES

b. **One** of the following:

- i. For treatment of familial chylomicronemia syndrome (FCS), has **one** of the following:
  - a) Results of genetic testing showing biallelic pathogenic variants in FCS-causing genes,
  - b) A North American FCS (NAFCS) score greater than or equal to 60 (i.e., definite FCS),
  - c) **Both** of the following:
    - (i) An NAFCS score greater than or equal to 45 and less than 60 (i.e., likely FCS)
    - (ii) An FCS score (Moulin score) greater than or equal to 10 (i.e., FCS very likely)
- ii. For all other diagnoses, has a history of therapeutic failure of or a contraindication or an intolerance to first line therapy(ies) if applicable according to consensus treatment guidelines;

**AND**

11. For all other non-preferred Lipotropics, Other, has a history of therapeutic failure of or a contraindication or an intolerance to the preferred Lipotropics, Other approved or medically accepted for the beneficiary's diagnosis; **AND**
12. If a prescription for a Lipotropics, Other is for a quantity that exceeds the quantity limit, the determination of whether the prescription is medically necessary will also take into account the guidelines set forth in the Quantity Limits Chapter.

NOTE: If the beneficiary does not meet the clinical review guidelines but, in the professional judgment of the physician reviewer, the services are medically necessary to meet the medical needs of the beneficiary, the request for prior authorization will be approved.

FOR RENEWALS OF PRIOR AUTHORIZATION FOR LIPOTROPICS, OTHER: The determination of medical necessity of a request for renewal of a prior authorization for a Lipotropics, Other that was previously approved will take into account whether the beneficiary:

1. Has documentation of a positive clinical response since starting the requested drug (e.g., decreased LDL-C, decreased triglycerides, fewer episodes of acute pancreatitis, etc.); **AND**
2. Is prescribed a dose that is consistent with FDA-approved package labeling, nationally recognized compendia, or peer-reviewed medical literature; **AND**

MEDICAL ASSISTANCE HANDBOOK  
PRIOR AUTHORIZATION OF PHARMACEUTICAL SERVICES

3. Does not have a contraindication to the prescribed drug; **AND**
4. For a PCSK9 inhibitor, is using the requested PCSK9 inhibitor in addition to **one** of the following:
  - a. For treatment of HoFH, standard lipid-lowering treatments as recommended by current consensus guidelines<sup>3</sup>
  - b. For treatment of all other conditions, the maximum tolerated dose of the highest-tolerated intensity statin (if clinically appropriate);

**AND**

5. For an ACL inhibitor, **both** of the following:
  - a. Is using the requested ACL inhibitor in addition to the maximum tolerated dose of the highest-tolerated intensity statin (if clinically appropriate)
  - b. If currently taking simvastatin or pravastatin, is not using the requested ACL inhibitor concomitantly with simvastatin at a dose of greater than 20 mg daily or pravastatin at a dose of greater than 40 mg daily;

**AND**

6. For an ANGPTL3 inhibitor or MTP inhibitor, **both** of the following:
  - a. Is prescribed the requested drug by or in consultation with a cardiologist, endocrinologist, or other provider specializing in lipid disorders
  - b. Is using the requested drug in addition to standard lipid-lowering treatments as recommended by current consensus guidelines;

**AND**

7. For icosapent ethyl, experienced a decrease in fasting triglycerides since starting icosapent ethyl; **AND**
8. For an *APOC-III*-directed ASO, is prescribed the requested drug by or in consultation with a cardiologist, endocrinologist, gastroenterologist, or other provider specializing in lipid disorders; **AND**
9. For all other non-preferred Lipotropics, Other, has a history of therapeutic failure of or a contraindication or an intolerance to the preferred Lipotropics, Other approved or medically accepted for the beneficiary's diagnosis; **AND**

---

<sup>3</sup> e.g., American Heart Association/American College of Cardiology, American Association of Clinical Endocrinologists/American College of Endocrinology, American Diabetes Association, National Lipid Association, European Society of Cardiology/European Atherosclerosis Society, International Familial Hypercholesterolaemia Foundation, International Atherosclerosis Society

MEDICAL ASSISTANCE HANDBOOK  
PRIOR AUTHORIZATION OF PHARMACEUTICAL SERVICES

10. If a prescription for a Lipotropics, Other is for a quantity that exceeds the quantity limit, the determination of whether the prescription is medically necessary will also take into account the guidelines set forth in the Quantity Limits Chapter.

NOTE: If the beneficiary does not meet the clinical review guidelines listed above but, in the professional judgment of the physician reviewer, the services are medically necessary to meet the medical needs of the beneficiary, the request for prior authorization will be approved.

C. Clinical Review Process

Prior authorization personnel will review the request for prior authorization and apply the clinical guidelines in Section B. above to assess the medical necessity of a prescription for a Lipotropics, Other. If the guidelines in Section B. are met, the reviewer will prior authorize the prescription. If the guidelines are not met, the prior authorization request will be referred to a physician reviewer for a medical necessity determination. Such a request for prior authorization will be approved when, in the professional judgment of the physician reviewer, the services are medically necessary to meet the medical needs of the beneficiary.

D. References

1. Evkeeza [package insert]. Tarrytown, NY: Regeneron Pharmaceuticals, Inc. April 2023.
2. Juxtapid [package insert]. Dublin, Ireland: Amryt Pharmaceuticals DAC. September 2020.
3. Leqvio [package insert]. East Hanover, NJ: Novartis Pharmaceuticals Corporation. June 2024.
4. Nexletol [package insert]. Ann Arbor, MI: Esperion Therapeutics, Inc. March 2024.
5. Nexlizet [package insert]. Ann Arbor, MI: Esperion Therapeutics, Inc. March 2024.
6. Praluent [package insert]. Tarrytown, NY: Regeneron Pharmaceuticals Inc.; March 2024.
7. Repatha [package insert]. Thousand Oaks, CA: Amgen Inc. September 2021.
8. Tryngolza [package insert]. Carlsbad, CA: Ionis Pharmaceuticals, Inc.; December 2024.
9. Jellinger PS, Handelsman Y, Rosenblit PD, et al. American Association of Clinical Endocrinologists and American College of Endocrinology guidelines for management of dyslipidemia and prevention of cardiovascular disease. *Endocri Pract.* 2017;23(Suppl. 2):1-87.
10. Grundy SM, Stone NJ, Bailey AL, et al. 2018 AHA/ACC/AACVPR/AAPA/ABC/ACPM/ADA/AGS/APhA/ASPC/NLA/PCNA guideline on the management of blood cholesterol: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *Circulation.* 2019;139:e1082-e1143.
11. The Task Force for the management of dyslipidaemias of the European Society of Cardiology (ESC) and European Atherosclerosis Society (EAS). 2019 ESC/EAS guidelines for the management of dyslipidaemias: lipid modification to reduce cardiovascular risk. *Eur Heart J.* 2019;00:1-78.
12. Kleindorfer DO, Towfighi A, Chaturvedi S, et al. 2021 guideline for the prevention of stroke in patients with stroke and transient ischemic attack: A guideline from the American Heart Association/American Stroke Association. *Stroke.* 2021;52:e364-e467.
13. Visseren JLF, Mach F, Smulders VM, et al. 2021 ESC guidelines on cardiovascular disease prevention in clinical practice. *Eur Heart J.* 2021;42:3227-3337.
14. Virani SS, Morris PB, Agarwala A, et al. 2021 ACC excerpt consensus decision pathway on the management of ASCVD risk reduction in patients with persistent hypertriglyceridemia: A report of the American College of Cardiology Solution Set Oversight Committee. *J Am Coll Cardiol.* 2021;78(9):960-993.
15. Lloyd-Jones DM, Morris PB, Ballantyne CM, et al. 2022 ACC expert consensus decision pathway role of nonstatin therapies for LDL-cholesterol lowering in the management of atherosclerotic cardiovascular disease

MEDICAL ASSISTANCE HANDBOOK  
PRIOR AUTHORIZATION OF PHARMACEUTICAL SERVICES

risk: A report of the American College of Cardiology Solution Set Oversight Committee. *J Am Coll Cardiol.* 2022.

16. Rosenson RS, Hayward RA, Lopez-Sendon J. Management of low density lipoprotein cholesterol (LDL-C) in the secondary prevention of cardiovascular disease. In: UpToDate [internet database]. Freeman MW, Cannon CP, Parikh N, eds. Waltham, MA: UpToDate Inc. Updated April 1, 2022. Accessed April 18, 2022.
17. Rosenson RS, Eckel RH. Hypertriglyceridemia in adults: Management. In: UpToDate [internet database]. Freeman MW, Parikh N, Givens J, eds. Waltham, MA: UpToDate Inc. Updated March 4, 2022. Accessed July 6, 2022.

Inherited Dyslipidemias

18. Gidding SS, Champagne MA, de Ferranti SD, et al. The agenda for familial hypercholesterolemia – a scientific statement from the American Heart Association. *Circulation.* 2015;132:2167-2192.
19. Santos RD, Gidding SS, Hegele RA, et al. Defining severe familial hypercholesterolemia and the implications for clinical management: a consensus statement from the International Atherosclerosis Society Severe Familial Hypercholesterolemia Panel. *Lancet Diabetes-Endocrinol.* 2016;4(10):850-861.
20. Wiegman A, Gidding SS, Watts GF, et al. Familial hypercholesterolaemia in children and adolescents: gaining decades of life by optimizing detection and treatment. *Eur Heart J.* 2015;36(36):2425-2437.
21. France M, Rees A, Datta D, et al. HEART UK statement on the management of homozygous familial hypercholesterolaemia in the United Kingdom. *Atherosclerosis.* 2016;255:128-139.
22. Cuchel M, Bruckert E, Ginsberg HN, et al. Homozygous familial hypercholesterolaemia: new insights and guidance for clinicians to improve detection and clinical management. A position paper from the Consensus Panel on Familial Hypercholesterolaemia of the European Atherosclerosis Society. *Eur Heart J.* 2014;35(32):2146-2157.
23. Raal FJ, Hovingh GK, Catapano AL. Familial hypercholesterolemia treatments: guidelines and new therapies. *Atherosclerosis.* 2018;277:483-492.
24. Hegele RA, Ahmad Z, Ashraf A, et al. Development and validation of clinical criteria to identify familial chylomicronemia syndrome (FCS) in North America. *J Clin Lipidol.* 2025;19:83-94.
25. Moulin P, Dufour R, Averna M, et al. Identification and diagnosis of patients with familial chylomicronaemia syndrome (FCS): Expert panel recommendations and proposal of an “FCS score”. *Atherosclerosis.* 2018;275:265-272.
26. Rosenson RS, Durrington P. Familial hypercholesterolemia in adults: treatment. In: UpToDate [internet database]. Freeman MW, Parikh N, eds. Waltham, MA: UpToDate Inc. Updated September 14, 2020. Accessed April 18, 2022.
27. Rosenson RS, Durrington P. Inherited disorders of LDL-cholesterol metabolism other than familial hypercholesterolemia. In: UpToDate [internet database]. Freeman MW, Cosentino F, Parikh N, eds. Waltham, MA: UpToDate Inc. Updated July 1, 2020. Accessed April 18, 2022.

Non-Statin Drugs

28. Landmesser U, Chapman MJ, Stock JK, et al. 2017 update of ESC/EAS Task Force on practical clinical guidance for proprotein convertase subtilisin/kexin type 9 inhibition in patients with atherosclerotic cardiovascular disease or in familial hypercholesterolaemia. *Eur Heart J.* 2018;39(14):1131-1143.
29. Lloyd-Jones DM, Morris PB, Ballantyne CM, et al. 2017 focused update of the 2016 ACC expert consensus decision pathway on the role of non-statin therapies for LDL-cholesterol lowering in the management of atherosclerotic cardiovascular disease risk: a report of the American College of Cardiology Task Force on Clinical Expert Consensus Decision Pathways. *J Am Coll Cardiol.* 2017;70(14):1785-1822.
30. Orringer CE, Jacobson TA, Saseen JJ, et al. Update on the use of PCSK9 inhibitors in adults: recommendations from an Expert Panel of the National Lipid Association. *J Clin Lipidol.* 2017;11:880-890.
31. Banach M, Penson PE, Farnier M, et al. Bempedoic acid in the management of lipid disorders and cardiovascular risk. 2023 position paper of the International Lipid Expert Panel (ILEP). *Prog Cardiovasc Dis.* 2023 Mar 7;S0033-0620(23)00026-9.

Statin Intolerance

32. Rosenson RS, Baker SK, Jacobson TA, et al. An assessment by the Statin Muscle Safety Task Force: 2014 update. *J Clin Lipidol.* 2014;8:S58-S71.
33. Bays H, Cohen DE, Chalasani N, Harrison SA. An assessment by the Statin Liver Safety Task Force: 2014 update. *J Clin Lipidol.* 2014;8:S47-S57.
34. Guyton JR, Bays HE, Grundy SM, Jacobson TA. An assessment by the Statin Intolerance Panel: 2014 update. *J Clin Lipidol.* 2014;8:S72-S81.

MEDICAL ASSISTANCE HANDBOOK  
PRIOR AUTHORIZATION OF PHARMACEUTICAL SERVICES

35. Stroes ES, Thompson PD, Corsini A, et al. Statin-associated muscle symptoms: impact on statin therapy – European Atherosclerosis Society Consensus Panel state on assessment, aetiology and management. *Eur Heart J*. 2015;36:1012-1022.
36. Banach M, Rizzo M, Toth P, et al. Statin intolerance – an attempt at a unified definition. Position paper from an International Lipid Expert Panel. *Arch Med Sci*. 2015;11(1):1-23.
37. Mancini GBJ, Baker S, Bergeron J, et al. Diagnosis, prevention, and management of statin adverse effects and intolerance: Canadian Consensus Working Group update (2016). *Can J Cardiol*. 2016;32:S35-S65.
38. Newman CB, Preiss D, Tobert JA, et al. Statin safety and associated adverse events: A scientific statement from the American Heart Association. *Arterioscler Thromb Vasc Biol*. 2019;39:e38-e81.
39. Cheeley MK, Saseen JJ, Agarwala A, et al. NLA scientific statement on statin intolerance: A new definition and key considerations for ASCVD risk reduction in the statin intolerant patient. *J Clin Lipidol*. 2022.
40. Rosenson RS, Baker SK. Statin myopathy. In: UpToDate [internet database]. Freeman MW, Rind DM, eds. Waltham, MA: UpToDate. Updated July 10, 2015.
41. Rosenson RS, Baker SK. Statin muscle-related adverse events. In: UpToDate [internet database]. Freeman MW, Givens J, eds. Waltham, MA: UpToDate Inc. Updated February 25, 2019. Accessed August 9, 2019.